

Interaction corrections at intermediate temperatures: dephasing time

B.N. Narozhny^(a), Gábor Zala^(b) and I.L. Aleiner^(b)

^(a)Physics Department, Brookhaven National Laboratory, Upton, NY, 11973-5000, USA

^(b)Department of Physics and Astronomy, SUNY at Stony Brook, Stony Brook, NY, 11794, USA
(February 1, 2008)

We calculate the temperature dependence of the weak localization correction in a two dimensional system at arbitrary relation between temperature, T and the elastic mean free time. We describe the crossover in the dephasing time $\tau_\varphi(T)$ between the high temperature, $1/\tau_\varphi \simeq T^2 \ln T$, and the low temperature $1/\tau_\varphi \simeq T$ behaviors. The prefactors in these dependences are not universal, but are determined by the Fermi liquid constant characterising the spin exchange interaction.

PACS numbers: 72.10.-d, 71.30.+h, 71.10.Ay

Introduction – The concept of a time scale characteristic of electron scattering processes in metals has been the focus of intense theoretical research for the last three decades^{1–16}. It has been established for disordered systems⁴ that the time scale, which corresponds to processes that suppress quantum interference (and are thus responsible for the temperature dependence of the weak localization correction, for example), namely the phase relaxation time τ_φ , is quite different from the semi-classical time scales, such as the energy relaxation time τ_E .

Previous work was mostly focused on the universal (independent of interaction strength) contribution of the singlet channel interaction both in the diffusive^{4,5} and ballistic^{3,5,11,13,14} regimes. The contribution of the triplet channel was only considered in the diffusive regime in Ref. 8. In this paper we fill the gaps by considering both channels at arbitrary relation between temperature and the inverse mean free time and thus describing the cross-over between the diffusive and ballistic regimes. We also clarify the relation between thus calculated dephasing time and experimentally observable physical quantities.

A discussion of phase relaxation should begin by defining a physical quantity sensitive to quantum interference (since the phase by itself is not an observable quantity). Therefore, the precise definition of the phase relaxation time depends on the choice of such physical quantity. Consider the weak localization correction in two dimensions^{6,15} in the absence of magnetic field:

$$\delta\sigma_{WL}(T) = -\frac{e^2}{2\pi^2\hbar} \int_{-\infty}^{\infty} \frac{d\epsilon}{4T \cosh^2 \frac{\epsilon}{2T}} \ln \frac{\tau_\varphi(\epsilon, T)}{\tau}, \quad (1)$$

where τ is the transport elastic mean free time. With the logarithmic accuracy one can here neglect the dependence of τ_φ on ϵ and estimate^{6,17}

$$\delta\sigma_{WL}(T) \approx -\frac{e^2}{2\pi^2\hbar} \ln \frac{\tau_\varphi(0, T)}{\tau}. \quad (2)$$

In the presence of magnetic field, H , perpendicular to the plane of the two-dimensional system, the situation

is more complicated. The magneto-conductivity can be described^{6,17,18} as

$$\sigma(H, T) - \sigma(0, T) = -\frac{e^2}{2\pi^2\hbar} \int_{-\infty}^{\infty} \frac{d\epsilon}{4T \cosh^2 \frac{\epsilon}{2T}} \times \left\{ Y[\Omega_H \tau_\varphi(\epsilon, T; H)] + \ln \left(\frac{\tau_\varphi(\epsilon, T; H)}{\tau_\varphi(\epsilon, T)} \right) \right\} \quad (3)$$

where $\Omega_H = 4DeH/\hbar c$, D is the diffusion constant, and

$$Y[x] = \ln \frac{1}{x} - \psi \left(\frac{1}{2} + \frac{1}{x} \right),$$

with $\psi(x)$ being the digamma function. Notice, that the dephasing time now depends on the magnetic field, see Refs. 6,18,15 and Eq. (8) below. We use the notation

$$\tau_\varphi(\epsilon, T) \equiv \tau_\varphi(\epsilon, T; H = 0).$$

One can simplify the magneto-conductivity (3) using asymptotic expressions for $Y[x]$ in order to facilitate comparison with experimental data. For strong magnetic fields $\Omega_H \tau_\varphi \gg 1$ one finds $Y[x \gg 1] \approx -\ln x + 2 \ln 2 + \mathbf{C}$, with $\mathbf{C} \approx 0.5772 \dots$ is the Euler constant. This yields

$$\sigma(H, T) - \sigma(0, T) = \frac{e^2}{2\pi^2\hbar} \ln \alpha \Omega_H \tau_\varphi^H + \mathcal{O} \left(\frac{1}{\Omega_H \tau_\varphi^H} \right) \quad (4)$$

where $\alpha = 1/(4e^{\mathbf{C}}) = 0.1403 \dots$, and

$$\tau_\varphi^H = \exp \left(\int_{-\infty}^{\infty} \frac{d\epsilon}{4T \cosh^2 \frac{\epsilon}{2T}} \ln \tau_\varphi(\epsilon, T) \right) \quad (5)$$

In the opposite limit of the weak field ($Y[x \ll 1] \approx -x^2/24$) the magneto-conductivity is quadratic in Ω_H

$$\sigma(H, T) - \sigma(0, T) = \frac{e^2}{48\pi^2\hbar} (\Omega_H \tau_\varphi^T)^2 \quad (6)$$

where

$$\tau_\varphi^T = \left[\int_{-\infty}^{\infty} \frac{d\epsilon}{4T \cosh^2 \frac{\epsilon}{2T}} \tau_\varphi^2(\epsilon, T) \right]^{1/2}. \quad (7)$$

As we show below, the two scales τ_φ^T and τ_φ^H may differ by a numerical factor which is important for a quantitative comparison with experiments.

As a function of ϵ , T and H the phase relaxation time is given by⁶

$$\frac{1}{\tau_\varphi(\epsilon, T; H)} = \int_{\max[\Omega_H, \tau_\varphi^{-1}]}^{\infty} d\omega \omega A(\omega) \times \left[\coth \frac{\omega}{2T} - \frac{1}{2} \left(\tanh \frac{\omega - \epsilon}{2T} + \tanh \frac{\omega + \epsilon}{2T} \right) \right]. \quad (8)$$

Kernel $A(\omega)$ contains all the information about matrix elements of the interaction and wavefunctions of the disordered system, see e.g.^{6,15}, and this is precisely the same kernel which enters into the inelastic collision integral in the Boltzmann equation, see e.g. Ref. 6,16:

$$\text{St}_{in}\{f(\epsilon)\} = \int d\omega \int d\epsilon_1 A(\omega) f(\epsilon_1) [1 - f(\epsilon_1 - \omega)] \times \left\{ -f(\epsilon) [1 - f(\epsilon + \omega)] + [1 - f(\epsilon)] f(\epsilon - \omega) \right\}, \quad (9)$$

where f is the distribution function of the quasiparticles.

Calculation of the kernel – The kernel $A(\omega)$ can be obtained from the quantum kinetic equation (see Ref. 19) and it is expressed in terms of the interaction propagator $\mathcal{D}^R$ and the propagators $\langle D \rangle$ describing semi-classical dynamics of non-interacting electrons

$$A(\omega) = \frac{2\nu}{\pi} \int \frac{d^2 q}{(2\pi)^2} \left([\mathbf{Re} \langle D \rangle]^2 |\mathcal{D}_S^R(\omega, q)|^2 + \text{Tr}[\mathbf{Re} \langle \hat{D} \rangle] \hat{\mathcal{D}}_T^R(\omega, q) [\mathbf{Re} \langle \hat{D} \rangle] [\hat{\mathcal{D}}_T^R(\omega, q)]^* \right). \quad (10)$$

In the absence of external magnetic field and spin-orbit interaction the semi-classical propagator is diagonal $\langle \hat{D} \rangle = \delta_{ij} \langle D \rangle$ and

$$\langle D \rangle = \frac{1}{\sqrt{(-i\omega + 1/\tau)^2 + v_F^2 q^2} - 1/\tau}. \quad (11)$$

It becomes the usual diffuson⁶ in the diffusive limit $\omega, v_F q \ll 1/\tau$.

The interaction propagator in the singlet channel is given in terms of $\langle D \rangle$ and the Fermi liquid constant $F_0^{\rho 19,20}$

$$\mathcal{D}_S^R(\omega, q) = -\frac{1}{\nu} \frac{\nu V_0(q) + F_0^\rho}{1 + (\nu V_0(q) + F_0^\rho)[1 + i\omega \langle D \rangle]}. \quad (12)$$

The interaction propagator in the triplet channel is a diagonal matrix in spin indices:

$$\hat{\mathcal{D}}_T^R(\omega, q) = -\frac{\delta_{ij}}{\nu} \frac{F_0^\sigma}{1 + F_0^\sigma [1 + i\omega \langle D \rangle]}, \quad (13)$$

where F_0^σ is the Fermi-liquid constant in the triplet channel^{19,20}. At distances larger than the screening radius one can take the unitary limit in Eq. (12), $\nu V_0(q) \rightarrow \infty$ which coincides with the $F_0^\sigma \rightarrow \infty$ limit in Eq. (13) [see Ref. 19 for more details].

We now evaluate the triplet contribution to the kernel $A(\omega)$, separating Eq. (10) into the sum

$$A(\omega) = 3A_T(\omega) + A_S(\omega). \quad (14)$$

Using Eqs. (13) and (11) one can rewrite Eq. (10) as

$$A_T(\omega) = \frac{1}{\pi\omega} \text{Im} \int \frac{d^2 q}{(2\pi)^2} \mathcal{D}_T^R(\omega, q) [\langle D \rangle + \langle D \rangle^*] \quad (15)$$

The kernel $A_S(\omega)$ can be obtained from Eq. (15) by taking the limit $F_0^\sigma \rightarrow \infty$. The momentum integral in Eq. (15) diverges logarithmically in the ultraviolet. This divergence is typically^{5,3,14} cut off at a scale of order k_F (the precise definition of such cut off is important only for the numerical factor under the logarithm which we believe is beyond the accuracy of any discussion). In the diffusive limit the divergence does not appear since one is limited by small momenta $v_F q \ll 1/\tau$. It is therefore convenient to separate $A_T(\omega)$ into two parts, roughly corresponding to the ballistic and diffusive asymptotics, respectively:

$$A_T(\omega) = A_1(\omega) + A_2(\omega). \quad (16)$$

The “ballistic” term appears from integrating over momenta larger than inverse elastic mean free time and/or frequency. With logarithmic accuracy we find (here the product of the Fermi momentum and the Fermi velocity [renormalized by interaction] is denoted by $E_F = v_F k_F/2$)

$$A_1(\omega) = \frac{(F_0^\sigma)^2}{4\pi E_F (1 + F_0^\sigma)^2} \ln \frac{E_F^2}{\tau^{-2} + b(F_0^\sigma)\omega^2}, \quad (17a)$$

$$b(x) \approx \frac{1 + x^2}{(1 + x)^2}$$

This term has the logarithmic frequency dependence similar to the one obtained by several authors for the singlet channel interaction^{5,14}.

The “diffusive” term, i.e. coming from integrating over small momenta, is a generalization of the standard result⁶. For simplicity, we show the kernel $A_2(\omega)$ in the two limiting cases. For small energy transfers $\omega\tau \ll 1$ we find

$$A_2(\omega\tau \ll 1) = \frac{F_0^\sigma}{(1 + F_0^\sigma)(2 + F_0^\sigma)} \frac{1}{2g\omega} \times \left[(1 + F_0^\sigma) \arctan \frac{1}{\omega\tau} - \arctan \frac{1 + F_0^\sigma}{\omega\tau} \right]. \quad (17b)$$

where $g = 2\pi\hbar/e^2 R_\square$ is the dimensionless conductance of the system, and $R_\square$ is the sheet resistance.

Note, that this result reduces to that of the diffusive theory for $\omega\tau \ll 1 + F_0^\sigma$ (see Refs. 19 and 21 for a more detailed discussion of the crossover to the ballistic regime). For the singlet channel (which can be obtained by setting $F_0^\sigma \rightarrow \infty$) Eq. (17b) coincides with the standard result⁶.

For large energy transfers $\omega\tau \gg 1$ integration over small moments yields a correction to Eq. (17a):

$$A_2(\omega\tau \gg 1) = \frac{1}{1 + F_0^\sigma} \frac{1}{2g\omega} \times \left[\arctan \frac{1 + F_0^\sigma}{\omega\tau(1 - F_0^\sigma)} + (1 + F_0^\sigma) \arctan \frac{1}{\omega\tau} \right]. \quad (17c)$$

For numerical reasons, contributions of Eq. (17c) to the final results are small compared with that of Eq. (17a) for all temperature regimes.

Results for the dephasing time – We now use the explicit form of the kernel Eqs. (17) to find the dephasing time from the self-consistency equation (8). According to Eqs. (14) and (16), it takes the form

$$\frac{1}{\tau_\varphi(\epsilon, T; H)} = I_1(\epsilon, T) + I_2(T, \max[\Omega_H, \tau_\varphi^{-1}]). \quad (18)$$

$$\frac{1}{\tau_\varphi^T} = \frac{1}{\tau_\varphi^H} = \left(1 + \frac{3(F_0^\sigma)^2}{(1 + F_0^\sigma)(2 + F_0^\sigma)} \right) \frac{T}{g} \ln [g(1 + F_0^\sigma)] + \frac{\pi}{4} \left(1 + \frac{3(F_0^\sigma)^2}{(1 + F_0^\sigma)^2} \right) \frac{T^2}{E_F} \ln(E_F \tau). \quad (20)$$

At stronger fields $\Omega_H > T$ equation (20) is not applicable, since in that regime the interaction correction in the Cooper channel becomes of the same order as the weak localization correction⁶. We will not dwell on this issue here.

The result (20) are valid while the condition $g(1 + F_0^\sigma) \gg 1$ holds. The same condition guarantees the exponential smallness of temperature independent dephasing induced by spontaneously spin polarized regions²².

Comparing the dominant, diffusive term [the first term in Eq. (20), which came from integrating Eq. (17b)] to the second, ballistic term, we find that the two become of the same order when $T\tau \sim 1 + F_0^\sigma$, which sets the limit of applicability to the purely diffusive theory.

2. Ballistic limit. At temperatures $T\tau \gg 1 + F_0^\sigma$ the leading asymptotics is controlled by Eq. (19a). In this regime we can no longer neglect the ϵ dependence in $\tau_\varphi(\epsilon, T)$. As a result, the time scales τ_φ^T , see Eq. (7), and τ_φ^H , see Eq. (5), are different by a numerical factor:

$$\tau_\varphi^{H,T} = \tau_\varphi(0, T) B^{H,T} \quad (21a)$$

where

The kernel (17a) is not diverging at small energy transfers, and the calculation of term I_1 can be performed by setting the lower limit of integration in Eq. (8) to zero. We find with the logarithmical accuracy

$$I_1 = \frac{\pi^2 T^2 + \epsilon^2}{8\pi E_F} \times \left[\frac{3(F_0^\sigma)^2}{(1 + F_0^\sigma)^2} \ln \left(\frac{E_F^2}{b(F_0^\sigma)T^2 + \tau^{-2}} \right) + \ln \left(\frac{E_F^2}{T^2 + \tau^{-2}} \right) \right], \quad (19a)$$

where function $b(x)$ is defined in Eq. (17a).

The diffusive term I_2 is logarithmically divergent at the lower limit. The divergent contribution is independent of ϵ and the divergence is cut according to Eq. (8)

$$I_2(T, \Omega) = \left(1 + \frac{3(F_0^\sigma)^2}{(1 + F_0^\sigma)(2 + F_0^\sigma)} \right) \frac{T}{g} \ln \left(\frac{T}{\Omega} \right). \quad (19b)$$

1. Diffusive limit ($T\tau \ll 1 + F_0^\sigma$). In this case the dephasing time follows from the self-consistency Eq. (8) where one has to include the singlet channel contribution. The time scales $\tau_\varphi^{T,H}$ defined in Eq. (4) are given by

$$\frac{1}{\tau_\varphi(0, T)} = \frac{\pi T^2}{4E_F} \left[\ln \frac{E_F}{T} + \frac{3(F_0^\sigma)^2}{(1 + F_0^\sigma)^2} \ln \frac{E_F}{T\sqrt{b(F_0^\sigma)}} \right], \quad (21b)$$

where $b(x)$ is defined in Eq. (17a), and the numerical factors are

$$B^T = \left[\int_0^\infty \frac{dz}{\cosh^2 z} \left(1 + \frac{4z^2}{\pi^2} \right)^{-2} \right]^{1/2} \approx 0.8437 \dots;$$

$$B^H = \exp \left[- \int_0^\infty \frac{dz}{\cosh^2 z} \ln \left(1 + \frac{4z^2}{\pi^2} \right) \right] \approx 0.7931 \dots$$

The observable dephasing times are different from $\tau_\varphi(0, T)$ by the above numerical constants. The temperature dependence of $\tau_\varphi(0, T)$ Eq. (21b) coincides with that found earlier in Refs. 3,5,11,13,14 for the case $F_0^\sigma = 0$ ^{23,24}. The numerical factor $\pi/4$ is the same as in Refs. 11,13,14, while Ref. 5 reports the factor $\pi/2$ and Ref. 3 reports the factor of $1/2\pi$. The correct numerical factor $\pi/4$ was also obtained in Ref. 12, however it was claimed that this result should be further renormalized (reduced by a factor of 4) by taking into

account higher order forward scattering processes. We believe such renormalization is erroneous and is a result of misidentification of the Fermi liquid constant in the singlet channel²⁵.

Summary – We have calculated the temperature dependence of the weak localization correction at arbitrary relation between T and elastic mean free time τ . The prefactors in the temperature dependencies of the dephasing rate are not universal and are determined by the single Fermi liquid constant F_0^σ . The very same constant determines the temperature dependence of the longitudinal resistivity¹⁹, the Hall coefficient²¹, and magnetoresistance in the parallel magnetic field²⁶. Because the number of different observable quantities exceeds the number of input parameters this theory possesses the predictive power.

Acknowledgments – We are grateful to M.Yu. Reizer and B. Laikhtman for interesting discussions, and to A.D. Mirlin for pointing an error in an earlier version of this paper. One of us (I.A.) was supported by the Packard foundation. Work at BNL is supported by the US DOE under Contract No DE-AC02-98CH10886.

-
- ¹ C. Hodges, H. Smith, and J.W. Wilkins, Phys. Rev. B **4**, 302 (1971).
 - ² A.V. Chaplik, Zh. Eksp. Teor. Fiz. **60**, 1845 (1971) [Sov. Phys. JETP **33**, 997 (1971)].
 - ³ G.F. Giuliani and J.J. Quinn, Phys. Rev. B **26**, 4421 (1982).
 - ⁴ B.L. Altshuler, A.G. Aronov, and D.E. Khmelnitsky, J. Phys. C **15**, 7367 (1982).
 - ⁵ H. Fukuyama and E. Abrahams, Phys. Rev. B **27**, 5976 (1983).
 - ⁶ B.L. Altshuler and A.G. Aronov in *Electron-Electron Interactions in Disordered Systems*, eds. A.L. Efros, M. Pollak (North-Holland, Amsterdam, 1985).
 - ⁷ P.A. Lee and Ramakrishnan, Rev. Mod. Phys. **57**, 287 (1985).
 - ⁸ C. Castellani, C. DiCastro, G. Kotliar, and P.A. Lee, Phys. Rev. Lett. **56**, 1179 (1986).
 - ⁹ H. Fukuyama and Y. Hasegawa, Prog. Theor. Phys. Suppl. **101**, 441 (1990).
 - ¹⁰ H. Fukuyama, Y. Hasegawa, and O. Narikiyo, J. Phys. Soc. Jpn. **60**, 1013 (1991).

- ¹¹ L. Zheng and S. Das Sarma, Phys. Rev. B **53**, 9964 (1996).
- ¹² D. Menashe and B. Laikhtman, Phys. Rev. B **54**, 11561 (1996).
- ¹³ T. Jungwirth and A.H. MacDonald, Phys. Rev. B **53**, 7403 (1996).
- ¹⁴ M. Reizer and J.W. Wilkins, Phys. Rev. B **55**, R7363 (1997).
- ¹⁵ I.L. Aleiner, B.L. Altshuler, and M.E. Gershenson, Waves Random Media **9**, 201 (1999).
- ¹⁶ I.L. Aleiner and Ya.M. Blanter, Phys. Rev. B **65**, 115317 (2002).
- ¹⁷ B.L. Altshuler, D. Khmelnitskii, A.I. Larkin, and P.A. Lee, Phys. Rev. B **22**, 5142 (1980); S. Hikami, A.I. Larkin, and Y. Nagaoka, Prog. Theor. Phys. **63**, 707 (1980).
- ¹⁸ E.L. Altshuler, B.L. Altshuler, and A.G. Aronov, Solid State Commun., **54**, 617 (1985).
- ¹⁹ Gábor Zala, B.N. Narozhny, and I.L. Aleiner, Phys. Rev. B **64**, 214204 (2001).
- ²⁰ L.D. Landau, Zh. Eksp. Teor. Fiz. **30**, 1058 (1956); *ibid* **32**, 59 (1957).
- ²¹ Gábor Zala, B.N. Narozhny, and I.L. Aleiner, Phys. Rev. B **64**, 201201 (2001).
- ²² B.N. Narozhny, I.L. Aleiner, and A.I. Larkin, Phys. Rev. B **62**, 14898 (2000).
- ²³ We note that the expression of the relaxation rate in terms of the Fermi liquid constant is somewhat similar to the taking into account of so called “local field corrections” of Ref. 13. We believe, that the deviation of our results from Eq. (24) of this reference is caused by the fact that the singlet (charge) and triplet, $L_z = 0$, contributions were not separated properly [see Eq. (19) of this reference].
- ²⁴ Reference 14 argues that the contribution of other “non-golden rule” diagrams, see Fig. 1 of Ref. 14 decreases the coefficient by a factor of 2. We think that this statement is not correct – this diagram, in fact, describes the renormalization of F_0^ρ and the effect of interaction in the Cooper channel on inelastic processes. The latter is suppressed by the factor of $1/\ln^2(E_F/T)$ for the repulsive interaction.
- ²⁵ The correct form of the propagator in the singlet channel is given by Eq. (12), which at distances greater than the screening radius reduces to $\mathcal{D}_S^R(q=0) = 1/\nu$. We believe that the resummation of Ref. 12 results in the incorrect form of the singlet propagator, $\mathcal{D}_S^R(q) = F_0^\rho/\nu + V(q)/[1 + \nu V(q)]$, where for forward scattering $F_0^\rho \rightarrow -1/2$, hence the extra factor of 1/4 in the final result of Ref. 12. Opinion of B. Laikhtman (private communication) is that disagreement by a factor of four is caused by algebraic mistakes in Eqs. (24) and (25) of Ref. 12.
- ²⁶ Gábor Zala, B.N. Narozhny, and I.L. Aleiner, Phys. Rev. B, **65**, 020201(R), (2002).